

The NuTeV Anomaly, Neutrino Mixing, and a Heavy Higgs

Tatsu Takeuchi

IPPAP, Physics Department, Virginia Tech, Blacksburg, VA 24061, U.S.A.

Talk presented at the YITP workshop ‘Progress in Particle Physics’, July 9, 2002.

The NuTeV experiment [1] has measured the ratios of neutral to charged current events in muon (anti)neutrino – nucleon scattering:

$$\begin{aligned} R_\nu &= \frac{\sigma(\nu_\mu N \rightarrow \nu_\mu X)}{\sigma(\nu_\mu N \rightarrow \mu^- X)} = g_L^2 + r g_R^2, \\ R_{\bar{\nu}} &= \frac{\sigma(\bar{\nu}_\mu N \rightarrow \bar{\nu}_\mu X)}{\sigma(\bar{\nu}_\mu N \rightarrow \mu^+ X)} = g_L^2 + \frac{g_R^2}{r}, \end{aligned} \quad (1)$$

where

$$r = \frac{\sigma(\bar{\nu}_\mu N \rightarrow \mu^+ X)}{\sigma(\nu_\mu N \rightarrow \mu^- X)} \sim \frac{1}{2}, \quad (2)$$

and has determined the parameters g_L^2 and g_R^2 to be

$$\begin{aligned} g_L^2 &= 0.3005 \pm 0.0014, \\ g_R^2 &= 0.0310 \pm 0.0011. \end{aligned} \quad (3)$$

The Standard Model (SM) predictions based on a fit to LEP/SLD data are cited as $[g_L^2]_{\text{SM}} = 0.3042$ and $[g_R^2]_{\text{SM}} = 0.0301$ in Ref. [1] and we see a 2.6σ disagreement in g_L^2 . In terms of the ratios R_ν and $R_{\bar{\nu}}$, this means that the neutral current events are not as numerous as the SM predicts when compared to the charged current events. On the LEP/SLD side, the measured invisible width is also known to be 2σ below the SM prediction. Both these facts seem to suggest that the $Z\nu\nu$ coupling is somehow suppressed.

Suppression of the $Z\nu\nu$ coupling can be arranged in a variety of models by mixing the neutrino with a heavy gauge singlet state as discussed in Ref. [3]. For the sake of simplicity, consider the case where all three generations of light neutrinos mix with a universal angle with such states. In such theories, if the $Z\nu\nu$ couplings are suppressed by a universal factor of $(1 - \varepsilon)$, then the $W\ell\nu$ couplings are also suppressed by a factor of $(1 - \varepsilon/2)$. Then the numerators of R_ν and $R_{\bar{\nu}}$ will be suppressed over their denominators, so such a mixing could in principle explain the NuTeV anomaly.

However, one must recall that one of the inputs used in calculating the SM predictions is the Fermi constant

G_F which is extracted from the muon decay constant G_μ . The suppression of the $W\ell\nu$ couplings would lead to the correction

$$G_F = G_\mu(1 + \varepsilon), \quad (4)$$

which will affect all SM predictions. The authors of Ref. [3] conclude that ε cannot be chosen to explain the NuTeV data without destroying the excellent agreement between LEP/SLD and the SM.

But what if we include oblique corrections from new physics [4] to compensate for the shift in G_F ? Note that the Fermi constant always appears multiplied by the ρ -parameter in neutral current amplitudes so a shift in G_F should be absorbable into the T -parameter.

In Ref. [5], we perform a global fit to the NuTeV and LEP/SLD data to determine what values of ε , S , and T are preferred. Using $m_H = 115$ GeV, $m_t = 174.3$ GeV as the reference SM, we obtain excellent agreement between theory and experiment at

$$\begin{aligned} S &= -0.05 \pm 0.10, \\ T &= -0.40 \pm 0.16, \\ \varepsilon &= 0.0028 \pm 0.0010. \end{aligned} \quad (5)$$

What kind of ‘new physics’ is compatible with these values of S and T ? It turns out that the simplest solution is the SM itself with a larger Higgs mass. Thus neutrino mixing together with a heavy Higgs will reconcile the LEP/SLD and NuTeV results.

Details of this analysis will be presented in Ref. [5].

Acknowledgments

I would like to thank Will Loinaz, Naotoshi Okamura, and L. C. R. Wijewardhana for their collaboration on this project, and Michio Hashimoto and Mihoko Nojiri for helpful discussions. This research was supported by the U.S. Department of Energy, grant DE-FG05-92ER40709, Task A.

-
- [1] G. P. Zeller *et al.* [NuTeV Collaboration], Phys. Rev. Lett. **88**, 091802 (2002) [hep-ex/0110059].
 - [2] The LEP Collaborations, the LEP Electroweak Working Group, and the SLD Heavy Flavor and Electroweak Groups, LEPEWWG/2002-01.
 - [3] S. Davidson, S. Forte, P. Gambino, N. Rius and A. Stru-

- mia, JHEP **0202**, 037 (2002) [hep-ph/0112302].
- [4] M. E. Peskin and T. Takeuchi, Phys. Rev. Lett. **65**, 964 (1990); Phys. Rev. **D46**, 381 (1992),
- [5] W. Loinaz, N. Okamura, T. Takeuchi, and L. C. R. Wijewardhana, in preparation.